

Product Information

CMV Immunoglobulin-VF

Australia

NAME OF THE MEDICINE

Human Cytomegalovirus Immunoglobulin, solution for intravenous injection.

DESCRIPTION

CMV Immunoglobulin-VF is a sterile, preservative free solution containing 55–65 mg/mL plasma proteins of which at least 98% is immunoglobulins (mainly IgG), with a cytomegalovirus (CMV) immunoglobulin activity of 1.5 million units per vial. The content of CMV immunoglobulin is defined in terms of EIA (enzyme immunoassay) units. One unit is equivalent to the specified volume of a standard CMV Immunoglobulin. Isotonicity is achieved by the addition of 292 mmol/L maltose. The distribution of the IgG subclasses closely resembles that found in normal human plasma (approximate mean ranges: 52.5–64.9% IgG₁, 29.0–42.2% IgG₂, 3.1–6.8% IgG₃, 0.4–1.2% IgG₄). CMV Immunoglobulin-VF contains only trace amounts of IgA (nominally <0.5 mg/mL).

CMV Immunoglobulin-VF is manufactured from human plasma collected by the Australian Red Cross Blood Service. Donations are selected on the basis that they contain high levels of specific antibodies to the CMV. The protein has not been chemically or enzymatically modified. However, the pH of the solution has been adjusted to 4.25. Immunoglobulins for intravenous injection, prepared by this process have not been implicated in the transmission of human immunodeficiency virus (HIV). Studies using plasma spiked with HIV have shown that the Cohn cold-ethanol fractionation process produces a very large reduction in virus titre with undetectable levels in the immunoglobulin fraction. Studies have shown that incubation at 27°C and pH 4.25, as performed in the manufacture of this product, also produces a very large reduction in HIV titre. Epidemiological studies have not recognised any cluster of AIDS patients or HIV seroconversion in immunoglobulin recipients. The manufacturing process for CMV Immunoglobulin-VF contains specific steps to reduce the possibility of viral transmission including low pH incubation for viral inactivation and nanofiltration for virus removal.

PHARMACOLOGY

The serum half-life of CMV Immunoglobulin-VF is normally approximately three weeks, but may be shortened considerably in patients who are immunosuppressed.

INDICATIONS

CMV Immunoglobulin-VF is indicated for the prevention of CMV infection following bone marrow and renal transplants. Specifically, the product is indicated when the recipient is seronegative for CMV and receives a graft from a CMV positive donor.

CMV Immunoglobulin-VF may also be a helpful adjunct to therapy in patients with established CMV infection, e.g. CMV pneumonitis.

CONTRAINDICATIONS

CMV Immunoglobulin-VF is contraindicated in individuals who have had a true anaphylactic reaction to a human immunoglobulin preparation. Individuals with isolated IgA deficiency should not receive the preparation, unless they have been tested and shown not to have circulating anti-IgA antibodies, since these patients may experience severe reactions to the IgA which is present in trace amounts.

PRECAUTIONS

CMV Immunoglobulin-VF MUST be administered intravenously only. Other routes of administration have not been evaluated. It is possible that CMV Immunoglobulin-VF may, on rare occasions, cause a precipitous fall in blood pressure and a clinical picture of anaphylaxis. These reactions may be related to the rate of infusion. Accordingly, the infusion rate given under **DOSAGE AND ADMINISTRATION** should be closely followed at least until the physician has had sufficient experience with a given patient. The patient's vital signs should be monitored regularly and careful observation made for any symptoms throughout the entire infusion. Adrenaline, oxygen, antihistamine and steroids should be available for the treatment of acute anaphylactic reactions.

An Aseptic Meningitis Syndrome (AMS) has been reported to occur infrequently in association with human intravenous immunoglobulin (IVIG) treatment. The syndrome usually begins within several hours to two days following IVIG treatment. It is characterised by symptoms and signs including severe headache, nuchal rigidity, drowsiness, fever, photophobia, painful eye movements, and nausea and vomiting. Cerebrospinal fluid (CSF) studies are frequently positive with pleocytosis, predominantly from the granulocytic series, and elevated protein levels. Patients exhibiting such symptoms and signs should receive a thorough neurological examination, including CSF studies, to rule out other causes of meningitis. AMS may occur more frequently in association with high dose (2 g/kg) IVIG treatment. Discontinuation of IVIG treatment has resulted in remission of AMS within several days without sequelae.

There have been occasional reports of renal dysfunction and acute renal failure in patients receiving IVIG products. Patients at increased risk are those with pre-existing renal insufficiency, diabetes mellitus, age greater than 65 years, volume depletion, sepsis and paraproteinaemia, and those taking concomitant nephrotoxic medicines. The majority of such incidents have been associated with sucrose-containing products. Whilst there is no sucrose in CMV Immunoglobulin-VF the following precautions should be followed: patients should be adequately hydrated prior to the initiation of the IVIG infusion and the recommended dose should not be exceeded. Renal function should be monitored in patients at increased risk of developing acute renal failure. If renal function deteriorates, discontinuation of IVIG should be considered.

There is clinical evidence of an association between IVIG administration and thromboembolic events which is assumed to be related to a relative increase in blood viscosity through the high influx of immunoglobulin in at-risk patients. Caution should be exercised when prescribing and infusing IVIG for patients with pre-existing risk factors for thrombotic events such as advanced age, estrogen use, in-dwelling vascular catheters, a history of venous or arterial thrombosis, acquired or inherited hypercoagulable states, cardiovascular risk factors (including history of atherosclerosis and/or impaired cardiac output), prolonged periods of immobilisation, severe hypovolaemia, hyperviscosity (including cryoglobulins, fasting chylomicronemia and/or high triglyceride levels, and monoclonal gammopathies). Reports have included cases of thrombophlebitis. In case of thromboembolic adverse reaction, the benefit and risk of treatment should be assessed before IVIG therapy is continued.

In patients at risk for thromboembolic adverse reactions, IVIG products should be administered at the minimum rate of infusion and dose practicable, and these individuals should be monitored for thrombotic complications. Consideration should also be given to measurement of baseline blood viscosity in individuals at risk for hyperviscosity.

IVIG products can contain blood group antibodies which may act as haemolysins and induce *in vivo* coating of red blood cells with immunoglobulin, causing a positive direct antiglobulin reaction (Coombs' test) and, rarely, haemolysis. Haemolytic anaemia can develop subsequent to IVIG therapy due to enhanced red blood cells (RBC) sequestration. Patients at increased risk for haemolysis following treatment with immunoglobulin include those with non-O blood group types, those who have underlying associated inflammatory conditions, and those receiving high cumulative doses of immunoglobulin over the course of several days. IVIG recipients should be monitored for clinical signs and symptoms of haemolysis, particularly

those patients at increased risk. If these occur, appropriate laboratory testing should be obtained.

In patients with limited or compromised acid-base compensatory mechanisms including neonates, consideration should be given to the effect of the additional acid load that the preparation might present.

Pathogen safety

This product is made from human plasma. Products made from human plasma may contain infectious agents, such as viruses and theoretically Creutzfeldt-Jakob Disease (CJD) agents that can cause disease. The risk that such products will transmit an infectious agent has been reduced by screening plasma donors for prior exposure to certain infectious agents and by testing for the presence of certain viral markers.

In addition, virus removal and inactivation steps are included in the manufacturing process. The current procedures applied in the manufacture of this product are effective against enveloped viruses such as human immunodeficiency virus (HIV), hepatitis B virus (HBV), hepatitis C virus (HCV), and the non-enveloped viruses, such as hepatitis A virus (HAV) and human parvovirus B19.

Despite these measures, such products may still potentially transmit disease. There is also the possibility that other known or unknown infectious agents may be present in such products.

Vaccination should be considered where appropriate, for patients in receipt of medicinal products manufactured from human plasma.

Genotoxicity, carcinogenicity and impairment of fertility

No genotoxicity, carcinogenicity or reproductive toxicity studies have been conducted with CMV Immunoglobulin-VF. There have been no reports of such effects associated with the use of CSL's plasma derived therapeutic medicines.

Use in pregnancy

The safety of CMV Immunoglobulin-VF for use in human pregnancy has not been established in controlled clinical trials and therefore it should be given to pregnant women only if clearly needed. However, clinical experience with other immunoglobulin preparations given during pregnancy suggests that there are no adverse effects on the foetus.

Use in lactation

Immunoglobulins are excreted in breast milk and may contribute to the transfer of protective antibodies to the neonate. It is not known, however, whether this applies to passively administered CMV Immunoglobulin-VF. Clinical experience with immunoglobulins suggests that no harmful effects on the neonate are to be expected.

Paediatric use and use in the elderly

The use of this product in the paediatric and elderly populations has not been established in appropriate studies. To date, these populations are not over-represented in spontaneous reports of adverse events associated with the use of CSL's IVIG therapeutic medicines.

Effect on laboratory tests

After injection of immunoglobulin, the transitory rise of the various passively transferred antibodies in the patient's blood may result in misleading positive results in serological testing.

Interference with glucose estimations

The maltose present in CMV Immunoglobulin-VF may interfere with some blood glucose measurements, resulting in the overestimation of blood glucose results. If this glucose measurement is used to guide treatment, hypoglycaemia may occur. Only certain glucose tests using glucose dehydrogenase have been implicated, so when monitoring glucose levels in patients receiving CMV Immunoglobulin-VF, information from the manufacturer of the glucose meter and/or test strips, should be reviewed to ensure that maltose does not interfere with the blood glucose reading.

INTERACTIONS WITH OTHER MEDICINES

Live attenuated virus vaccines: Passively acquired antibody can interfere with the response to live, attenuated virus vaccines. Therefore, administration of such vaccines, e.g. poliomyelitis or measles, should be deferred until approximately three months after passive immunisation. By the same token, immunoglobulins should not be administered for at least two weeks after a vaccine has been given.

ADVERSE EFFECTS

Reactions to IVIG tend to be related to the infusion rate and are most likely to occur during the first hour of the infusion. It is recommended that the patient's vital signs and general status be monitored regularly throughout the infusion.

The types of reactions that may occur include: abdominal pain, headache, chest-tightness, facial flushing or pallor, feeling hot, dyspnoea, non-urticarial skin rash, itching, hypotension, nausea or vomiting. Should any of these reactions develop during infusion of CMV Immunoglobulin-VF, the infusion should be temporarily stopped until the patient improves clinically (5 to 10 minutes) and then cautiously recommenced at a slower rate.

Some patients may develop delayed adverse reactions to CMV Immunoglobulin-VF such as nausea, vomiting, chest pain, rigors or aching legs. These adverse reactions occur after the infusion has stopped but usually within 24 hours.

True anaphylactic reactions to IVIG such as urticaria, angioedema, bronchospasm or hypotension occur very rarely. Should an anaphylactic reaction to CMV Immunoglobulin-VF develop, the infusion should be stopped and treatment instituted with adrenaline, oxygen, antihistamine and steroids.

There have been reports that IVIG can affect renal function. This should be monitored in patients with pre-existing renal failure who are given CMV Immunoglobulin-VF.

Rarely, renal dysfunction and acute renal failure have been reported.

DOSAGE AND ADMINISTRATION

Dosage

Because there is currently no international standard for CMV Immunoglobulin-VF, recommended dosage remains empirical. For prophylactic use it is recommended that patients receive 25,000 units/kg on days -4, -2, on the day of transplantation (intra-operatively) and then weekly for a further two months.

For therapy, 50,000 units/kg should be given initially, repeated after 4 to 5 days and then every 10 to 14 days until clinical improvement occurs.

Refer to **Table 1** for dosing requirements. If the calculated quantity of CMV Immunoglobulin-VF required for patients weighing >15 kg includes a fraction of a vial, then the higher whole number of vials should be administered.

Specifically, patients weighing ≤ 15 kg must be dosed based on their body weight (units/kg). Each vial of CMV Immunoglobulin-VF contains at least 20,000 units/mL of CMV immunoglobulin activity. The actual volume is stated on the label. If only a portion of the vial is used, discard the residual volume. Do not administer the whole vial. The volume and acid load should be considered if CMV Immunoglobulin-VF is administered to neonates.

Table 1: CMV Immunoglobulin-VF dosage based on body weight

Body weight (kg)	Dosage in number of vials	
	Prophylaxis	Therapy
0–≤15	<1 vial – Dosage based on body weight (units/kg)	
>15–30	1	1
31–60	1	2
61–90	2	3
91–120	2	4

Administration

If the product appears to be turbid by transmitted light or contains any sediment it must not be used, and the vial should be returned to the Australian Red Cross Blood Service. **The product does not contain an antimicrobial preservative. It must, therefore, be used immediately after opening the vial. Any unused solution must be discarded appropriately. Use in one patient on one occasion only. Do not use if the solution has been frozen.**

CMV Immunoglobulin-VF must be administered intravenously only.

CMV Immunoglobulin-VF may be infused undiluted. Allow the preparation to reach room temperature before infusing into the patient. Remove the dust cover from the top of the vial. **Apply a suitable antiseptic such as povidone-iodine or 70% ethanol to the exposed part of the rubber stopper and allow to dry.**

It is usual to gradually increase the infusion rate (provided the patient's vital signs are satisfactory at the lower rate) to the desired rate over the first 30 minutes. The infusion may be commenced at the rate of 1 mL per minute. After 15 minutes the rate may be gradually increased to a maximum of 3 to 4 mL per minute over a further 15 minutes.

Compatibility with other medicines

CMV Immunoglobulin-VF may be diluted with up to four times its volume of 0.9% saline or 5% glucose. No other medicine interactions or compatibilities have been evaluated.

CMV Immunoglobulin-VF should be administered separately from other intravenous fluids or medications the patient is receiving.

OVERDOSAGE

Overdosage may lead to fluid overload and hyperviscosity, particularly in the elderly and in patients with renal impairment.

PRESENTATION AND STORAGE CONDITIONS

CMV Immunoglobulin-VF is available in single vials each containing 1.5 million units of CMV immunoglobulin activity.

Store at 2°C to 8°C (Refrigerate. Do not freeze). Protect from light.

Do not use after the expiry date.

NAME AND ADDRESS OF THE SPONSOR

CSL Behring (Australia) Pty Ltd
ABN 48 160 734 761
189–209 Camp Road
Broadmeadows VIC 3047
Australia

DISTRIBUTED BY

Australian Red Cross Blood Service

POISON SCHEDULE OF THE MEDICINE

S4

DATE OF FIRST INCLUSION IN THE AUSTRALIAN REGISTER OF THERAPEUTIC GOODS (THE ARTG)

04 November 1991

DATE OF MOST RECENT AMENDMENT

15 October 2014

For Medical/Technical Enquiries

TOLL FREE: 1800 642 865

For Customer Service Enquiries

TOLL FREE: 1800 063 892

customerservice@cslbehring.com.au

www.cslbehring.com.au